

# $\mathcal{PT}$ symmetry and supersymmetry

**M Znojil**

Nuclear Physics Institute, 250 68 Řež, Czech Republic

**Abstract.** Pseudo-Hermitian (so called  $\mathcal{PT}$  symmetric) Hamiltonians are featured within a re-formulated Witten's supersymmetric quantum mechanics. An unusual form of the supersymmetric partnership between the spiked harmonic oscillators is described.

## 1. Supersymmetry for pedestrians, or bosons and fermions in Witten's picture

In the review [1] of the Witten's (or so called supersymmetric) quantum mechanics one finds the specific Fock vacuum

$$\langle q|0\rangle = \begin{bmatrix} \exp(-q^2/2)/\sqrt{\pi} \\ 0 \end{bmatrix}, \quad q \in (-\infty, \infty). \quad (1)$$

In the upper line we recognize the ground state of the one-dimensional harmonic oscillator  $H^{(HO)} = p^2 + q^2$  and see that “bosons” may be created/annihilated by the action of the respective operators  $-\partial_r + q$  and  $\partial_r + q$  (to be denoted as  $B^{(-1/2)}$  and  $A^{(-1/2)}$  here). The solvable harmonic-oscillator character of this elementary model enables us to define the three two-by-two matrices

$$\mathcal{F}^\dagger = \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}, \quad \mathcal{F} = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}, \quad \mathcal{N}_{\mathcal{F}} = \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \quad (2)$$

which create and annihilate a “fermion” and determine the fermionic number, respectively. Formally, this enables us to introduce the two partner Hamiltonians  $H_{(L)} = H^{(HO)} - 1$ ,  $H_{(R)} = H^{(HO)} + 1$  and identify the underlying symmetry with the superalgebra  $sl(1/1)$  generated by the following three operator matrices

$$\mathcal{H} = \begin{bmatrix} H_{(L)} & 0 \\ 0 & H_{(R)} \end{bmatrix}, \quad \mathcal{Q} = \begin{bmatrix} 0 & 0 \\ A & 0 \end{bmatrix}, \quad \tilde{\mathcal{Q}} = \begin{bmatrix} 0 & B \\ 0 & 0 \end{bmatrix}. \quad (3)$$

One easily verifies that  $\{\mathcal{Q}, \tilde{\mathcal{Q}}\} = \mathcal{H}$  while  $\{\mathcal{Q}, \mathcal{Q}\} = \{\tilde{\mathcal{Q}}, \tilde{\mathcal{Q}}\} = 0$  and  $[\mathcal{H}, \mathcal{Q}] = [\mathcal{H}, \tilde{\mathcal{Q}}] = 0$ .

Our forthcoming considerations may be summarized as a generalization of the above harmonic-oscillator-based model to  $D$  dimensions. Beyond its obvious phenomenological and methodical appeal, the mathematical motivation for such a construction stems from the well known requirements of absence of the centrifugal-type singularities in the general Witten's formalism [1]. Indeed, within the standard, Hermitian quantum mechanics, all ways of suppression of this difficulty seem to remain unclear up to these days [2]. In contrast, the weakening of the Hermiticity (to the so called  $\mathcal{PT}$  symmetry – see below) appears to be amazingly efficient in this context [3, 4].

## 2. $\mathcal{PT}$ symmetry for pedestrians, interpreted as a regularization of a spike in the force

Our key idea dates back to Buslaev and Grecchi [5] who proposed a specific regularization of the non-vanishing centrifugal term in  $D \neq 1$  dimensions (in their case, for some specific anharmonic oscillator examples) via a constant complex shift of the coordinate  $r = |\vec{q}|$ , better understood as a transition to the (increasingly popular [6]) non-Hermitian formalism of the so called  $\mathcal{PT}$  symmetric quantum mechanics [7]. In its present implementation, this merely means an extension of the real-line symmetry  $\mathcal{P}r = -r$  in (2.1) to the complex plane of  $r \in \mathbb{C}$ . Thus, we require the invariance of our Hamiltonian with respect to the parity  $\mathcal{P}$  multiplied by the time reversal mimicked by the complex conjugation,  $\mathcal{T}i = -i$ .

### 2.1. Illustration: $\mathcal{PT}$ symmetric version of the $D$ -dimensional harmonic oscillator

For the sake of brevity we shall only assign here our supersymmetry generators to the generalized class of the spiked harmonic oscillator Hamiltonians

$$H^{(\alpha)} = -\frac{d^2}{dr^2} + \frac{\alpha^2 - 1/4}{r^2} + r^2, \quad \alpha > 0.$$

Their Buslaev's and Grecchi's regularization will use  $r = x - i\varepsilon$  with real  $x$  and has thoroughly been studied in ref. [8]. Its normalizable wave functions

$$\psi(r) \equiv \mathcal{L}_N^{(\varrho)} = \langle r | N, \varrho \rangle = \frac{N!}{\Gamma(N + \varrho + 1)} \cdot r^{\varrho+1/2} \exp(-r^2/2) \cdot L_N^{(\varrho)}(r^2) \quad (4)$$

and energies

$$E = E_N^{(\varrho)} = 4N + 2\varrho + 2, \quad \varrho = -Q \cdot \alpha$$

are both labeled by an additional quantum number  $Q = \pm 1$  of the so called quasi-parity. In the context of fields, this concept proves closely related to the well known charge-conjugation symmetry  $\mathcal{C}$  [9].

## 3. $\mathcal{PT}$ symmetric supersymmetry and the pedestrian's spiked harmonic oscillator

We noticed in ref. [4] that the complexification of  $r$  regularizes the Witten's spiked harmonic oscillator (SHO) superpotential

$$W^{(\gamma)}(r) = -\frac{\partial_r \langle r | 0, \gamma \rangle}{\langle r | 0, \gamma \rangle} = r - \frac{\gamma + 1/2}{r} \quad r = r(x) = x - i\varepsilon, \quad x \in \mathbb{R}$$

as well as all the related operator matrix elements in the  $sl(1/1)$  generators (3),

$$\begin{aligned} A^{(\gamma)} &= \partial_r + W^{(\gamma)}, & B^{(\gamma)} &= -\partial_r + W^{(\gamma)}, & \gamma &\neq 0, \pm 1, \dots, \\ H_{(L)} &= B \cdot A = \hat{p}^2 + W^2 - W', & H_{(R)} &= A \cdot B = \hat{p}^2 + W^2 + W'. \end{aligned} \quad (5)$$

This returns us to the  $D = 1$  oscillator of section 1 at the “exceptional” value of  $\gamma = -1/2$  while at all the complex  $\gamma$  we have the generalized SUSY partners,

$$H_{(L)}^{(\gamma)} = H^{(\alpha)} - 2\gamma - 2, \quad H_{(R)}^{(\gamma)} = H^{(\beta)} - 2\gamma, \quad \alpha = |\gamma|, \quad \beta = |\gamma + 1|.$$

Whenever  $\gamma \in (-\infty, \infty)$  is real (while  $\alpha = |\gamma|$  and  $\beta = |\gamma + 1|$  are defined as positive), we have to distinguish between the following three different SUSY regimes characterized by the unbroken  $\mathcal{PT}$  symmetry,

$$\begin{cases} I. & \text{large negative } \gamma = -\alpha < -1, & \text{dominant } \alpha = \beta + 1 \\ II. & \text{small negative } \gamma = -\alpha > -1, & \text{both small, } \alpha + \beta = 1 \\ III. & \text{positive } \gamma = \alpha > 0, & \text{dominant } \beta = \alpha + 1 \end{cases}.$$

The energies (arranged in the descending order) form a (once degenerate) *completely real* quadruplet at each  $N$ ,

| SUSY partner energies | $I.$           | $II.$          | $III.$         |
|-----------------------|----------------|----------------|----------------|
| $E_{(L)}^{(\beta)}$   | $4N + 4\alpha$ | $4N + 4$       | $4N + 4$       |
| $E_{(L)}^{(\alpha)}$  | $4N + 4\alpha$ | $4N + 4\alpha$ | $4N$           |
| $E_{(L)}^{(-\beta)}$  | $4N + 4$       | $4N + 4\alpha$ | $4N - 4\alpha$ |
| $E_{(L)}^{(-\alpha)}$ | $4N$           | $4N$           | $4N - 4\alpha$ |

It is amusing to notice that up to the regular case (with  $\alpha = 1/2$ ) there always exist two alternative  $\gamma = \pm\alpha$  to a given  $\alpha > 0$ . Thus, each  $\alpha$  also has the *two different* partners  $\beta$  such that  $\beta_1 = |\alpha - 1| < \alpha < \beta_2 = \alpha + 1$ . Finally, in the domain II, all our SUSY construction remains perfectly valid even in the Hermitian limit  $\varepsilon \rightarrow 0$  [10].

#### 4. Non-standard $\mathcal{PT}$ symmetric supersymmetries

##### 4.1. Working at a fixed parameter $\alpha$

The respective annihilation and creation of ref. [4] was mediated by the *second-order* differential operators

$$A^{(-\gamma-1)} \cdot A^{(\gamma)} = A^{(\gamma-1)} \cdot A^{(-\gamma)} = \mathbf{A}(\alpha)$$

$$B^{(-\gamma)} \cdot B^{(\gamma-1)} = B^{(\gamma)} \cdot B^{(-\gamma-1)} = \mathbf{B}(\alpha)$$

with the “norm”  $c_5(N, \gamma) = -4\sqrt{(N+1)(N+\gamma+1)}$  and property

$$\mathbf{A}(\alpha) \cdot \mathcal{L}_{N+1}^{(\gamma)} = c_5(N, \gamma) \mathcal{L}_N^{(\gamma)}, \quad \mathbf{B}(\alpha) \cdot \mathcal{L}_N^{(\gamma)} = c_5(N, \gamma) \mathcal{L}_{N+1}^{(\gamma)}.$$

Hamiltonian  $H^{(\alpha)} = [\mathbf{A}(\alpha) \mathbf{B}(\alpha) - \mathbf{B}(\alpha) \mathbf{A}(\alpha)]/8$  satisfies commutation relations

$$\mathbf{A}(\alpha) H^{(\alpha)} - H^{(\alpha)} \mathbf{A}(\alpha) \equiv 4 \mathbf{A}(\alpha), \quad H^{(\alpha)} \mathbf{B}(\alpha) - \mathbf{B}(\alpha) H^{(\alpha)} \equiv 4 \mathbf{B}(\alpha)$$

of the Lie algebra  $sl(2, \mathbb{R})$  with the normalized generators  $\mathbf{A}(\alpha)/\sqrt{32}$ ,  $\mathbf{B}(\alpha)/\sqrt{32}$  and  $H^{(\alpha)}/4$ . As a consequence, the new,  $\mathcal{PT}$  SUSY results from eq. (3), with  $A$ ,  $B$  and  $H_{(L/R)}$  replaced by  $\mathbf{A}(\alpha)$ ,  $\mathbf{B}(\alpha)$  and  $\mathbf{G}_{(L/R)} = (\mathbf{H}^{(\alpha)} \mp 2)^2 - 4\alpha^2$ , respectively. The SHO eigenvectors themselves may then be obtained as solutions of the differential equations of the fourth order (cf. ref. [11]) which, in our case, read

$$\mathbf{G}_{(L)} |N^{(\gamma)}\rangle = \Omega_N^{(\gamma)} |N^{(\gamma)}\rangle, \quad \mathbf{G}_{(R)} |N^{(\gamma)}\rangle = \Omega_{N+1}^{(\gamma)} |N^{(\gamma)}\rangle.$$

where  $\Omega_N^{(\gamma)} = 16 N (N + \gamma)$ . This is our present main result.

#### 4.2. SUSY constructions at the complex $\gamma$

Marginally, let us note that even the complex choice of  $\gamma$  (when the  $\mathcal{PT}$  symmetry itself is broken) may lead to the partially real SUSY spectrum of energies. In order to show that, one has to derive a few identities for the Laguerre polynomials in (4) showing that the operators (5) change merely the subscripts or superscripts [4]. In the regime with the spontaneously broken  $\mathcal{PT}$  symmetry we may distinguish between the two options,

$$\begin{cases} \delta > 0 \text{ in } \gamma = i\delta, & \alpha = i\delta, & \beta = 1 + \alpha \\ \eta > 0 \text{ in } \gamma = -i\eta, & \alpha = i\eta, & \beta = 1 - \alpha \end{cases}$$

and get the partially real energy multiplets

$$\begin{cases} E_{(L)}^{(+\alpha)} = 4n, & E_{(L)}^{(-\alpha)} = E_{(R)}^{(-\beta)} = 4n - 4\alpha, & E_{(R)}^{(+\beta)} = 4n + 4 \\ E_{(L)}^{(-\alpha)} = 4n, & E_{(L)}^{(+\alpha)} = E_{(R)}^{(-\beta)} = 4n + 4\alpha, & E_{(R)}^{(+\beta)} = 4n + 4 \end{cases}$$

Similarly, at  $\gamma = N + i q \delta$  with  $q = \pm 1$  and  $\delta > 0$ , i.e., with no  $\mathcal{PT}$  symmetry at all, we get

$$E_{(L)} = \begin{cases} 4n - 4\gamma \\ 4n \end{cases}, \quad E_{(R)} = \begin{cases} 4n - 4\gamma \\ 4n + 4 \end{cases}$$

for the indices  $\beta_1 = N - 1 + i\delta$ ,  $\alpha = N + i\delta$  and  $\beta_2 = N + 1 + i\delta$ , still giving the partially real energy spectra.

#### Acknowledgements

Work partially supported by the grant of GA AS CR Nr. A 1048004.

#### References

- [1] Cooper F, Khare A and Sukhatme U 1995 Phys. Rep. 251 267
- [2] Jevicki A and Rodriguez J 1984 Phys. Lett. B 146 55;  
Das A and Pernice S 1999 Nucl. Phys. B 561 357;  
Gangopadhyaya A and Mallow J W 2002 ‘‘Supersymmetry in the Half-Oscillator - Revisited’’, LANL arXiv: hep-th/0206133;  
Das A and Pernice S 2002 ‘‘Comment on ‘Supersymmetry in the half-oscillator revisited’’’, LANL arXiv: hep-th/0207112
- [3] Znojil M 2000 Annihilation and creation operators in non-Hermitian supersymmetric quantum mechanics, arXiv hep-th/0012002
- [4] Znojil M 2002 J. Phys. A: Math. Gen. 35 2341
- [5] Buslaev V and Grecchi V 1993 J. Phys. A: Math. Gen. 26 5541
- [6] Bender C M or L  vai G or Mostafazadeh A or Quesne C 2002 Group 24 conference proceedings
- [7] Bender C M, Boettcher S and Meisinger P N 1999 J. Math. Phys. 40 2201
- [8] Znojil M 1999 Phys. Lett. A 259 220
- [9] Streater R F and Wightman A S 1964 PCT, spin and statistics and all that (New York: Benjamin);  
Bender C M, Brody D C and Jones H F 2002 Complex extension of quantum mechanics, arXiv: quant-ph/0208076
- [10] Znojil M 2002 Re-establishing supersymmetry between harmonic oscillators in  $D \neq 1$  dimensions, arXiv: hep-th/0203252
- [11] Bagchi B, Mallik S and Quesne C 2002 Int. J. Mod. Phys. A 17 51